

Experienced specialists. Advanced capabilities. Exceptional care



Experienced specialists. Advanced capabilities. Exceptional care.

Preparing for the Bravo™ pH Test with Colonoscopy

7 Days Before Your Test Start Date:

STOP all proton pump inhibitors (PPI) such as pantoprazole (Protonix[®]), omeprazole (Prilosec[®], Zegerid[®]), rabeprazole (AcipHex[®]), or lansoprazole (Prevacid[®]).

3 Days Before Your Test Start Date:

STOP H2 blockers such as ranitidine (Zantac®), cimetidine (Tagamet®), famotidine (Pepcid®, Fluxid®) or nizatidine (Axid®).

Purchase the Following Products:

- 1. Dulcolax Tablets (bisacodyl laxative)
- **2. Miralax 238 gram bottle** (to be mixed in the clear liquid choice below prior to drinking)
- 3. **Gatorade 64 ounces (NOTHING RED OR PURPLE)** Sugar-free drinks (G2, PowerAde Zero, Propel or water) are also options.

The Day BEFORE Your Exam:

- STOP taking over-the-counter antacids of any kind (Tums®, Rolaids®)
- STOP all solid foods. NO SOLID FOOD until after your exam.
- Beginning with a clear liquid breakfast, continue to drink at least 8 glasses (64 oz) of <u>CLEAR</u> liquids throughout the day and evening.
 Clear liquids include: Water, sports drinks, black coffee, tea, any soda (light or dark), broth, Jell-O, popsicles, and juice without pulp.
 NOTHING RED OR PURPLE.
- At 4:00 PM take four (4) of the Dulcolax Tablets.
- At <u>5:30 PM</u> drink half of the total Miralax/Gatorade solution over the next 2 hours.
- Continue drinking additional clear liquids throughout the evening.
- A soothing ointment (Desitin, Vaseline) may be applied to the anal area to prevent discomfort and skin breakdown.

The Day Of Your Exam:

• Four (4) hours before your arrival time, drink the remaining half of the Miralax/Gatorade solution over the next 2 hours.

2 Hours Before Your Arrival Time:

NOTHING BY MOUTH until after your test. No water, no gum, no mints. NOTHING BY MOUTH.

IF YOU ARE TO BE SEDATED, A
RESPONSIBLE DRIVER MUST REMAIN IN
THE FACILITY THROUGHOUT YOUR
APPOINTMENT. (~2 hours total)

Medications:

 Take all your regularly prescribed medications at least two (2) hours before you arrive.

Diabetics:

- Test your blood sugar before coming in for your procedure and bring the reading with you.
- If you take oral medication or insulin,
 DO NOT take it the morning of your procedure..
- Please contact the physician that manages your diabetic care if you have any questions regarding your oral medication or insulin dosages.

Blood Thinners:

- If you are taking blood thinners, call your prescribing physician and ask if you may stop taking the medication for the required amount of time prior to your procedure.
- You may continue taking aspirin, ibuprofen and other anti-inflammatory medications.

Bring with you:

- Your inhaler, if you use one and insulin, if you use it
- Completed registration forms if you did NOT use online registration
- Photo ID and Insurance Card
- List of all medications, vitamins and supplements you take (include dosage and last date and time each medication was taken before your arrival)
- Wear a short-sleeved shirt

Arrive at the <u>Endoscopy Center</u> at your scheduled time. Please call us if you have any questions regarding your preparation.

Granite Peaks Endoscopy 10150 S Petunia Way, Sandy, UT 84092 801-619-1115